



Bright Futures Parent Handout 1 Month Visit

Here are some suggestions from Bright Futures experts that may be of value to your family.

PARENTAL WELL-BEING

How You Are Feeling

- Taking care of yourself gives you the energy to care for your baby. Remember to go for your postpartum checkup.
- Call for help if you feel sad or blue, or very tired for more than a few days.
- Know that returning to work or school is hard for many parents.
- Find safe, loving child care for your baby. You can ask us for help.
- If you plan to go back to work or school, start thinking about how you can keep breastfeeding.

SAFETY

Safety

- Use a rear-facing car safety seat in all vehicles.
- Never put your baby in the front seat of a vehicle with a passenger air bag.
- Always wear your seat belt and never drive after using alcohol or drugs.
- Keep your car and home smoke-free.
- Keep hanging cords or strings away from and necklaces and bracelets off of your baby.
- Keep a hand on your baby when changing clothes or the diaper.

INFANT ADJUSTMENT

Getting to Know Your Baby

- Have simple routines each day for bathing, feeding, sleeping, and playing.
- Put your baby to sleep on his back.
 - In a crib, in your room, not in your bed.
 - In a crib that meets current safety standards, with no drop-side rail and slats no more than 2³/₈ inches apart. Find more information on the Consumer Product Safety Commission Web site at www.cpsc.gov.
 - If your crib has a drop-side rail, keep it up and locked at all times. Contact the crib company to see if there is a device to keep the drop-side rail from falling down.
 - Keep soft objects and loose bedding such as comforters, pillows, bumper pads, and toys out of the crib.
 - Give your baby a pacifier if he wants it.
- Hold and cuddle your baby often.
 - Tummy time—put your baby on his tummy when awake and you are there to watch.
- Crying is normal and may increase when your baby is 6–8 weeks old.
- When your baby is crying, comfort him by talking, patting, stroking, and rocking.
- *Never shake your baby.*
- If you feel upset, put your baby in a safe place; call for help.

FAMILY ADJUSTMENT

Your Baby and Family

- Plan with your partner, friends, and family to have time for yourself.
- Take time with your partner too.
- Let us know if you are having any problems and cannot make ends meet. There are resources in our community that can help you.
- Join a new parents group or call us for help to connect to others if you feel alone and lonely.
- Call for help if you are ever hit or hurt by someone and if you and your baby are not safe at home.
- Prepare for an emergency/illness.
 - Keep a first-aid kit in your home.
 - Learn infant CPR.
 - Have a list of emergency phone numbers.
 - Know how to take your baby's temperature rectally. Call us if it is 100.4°F (38.0°C) or higher.
- Wash your hands often to help your baby stay healthy.

Feeding Your Baby

- Feed your baby only breast milk or iron-fortified formula in the first 4–6 months.

FEEDING ROUTINES

- Pat, rock, undress, or change the diaper to wake your baby to feed.
- Feed your baby when you see signs of hunger.
 - Putting hand to mouth
 - Sucking, rooting, and fussing
- End feeding when you see signs your baby is full.
 - Turning away
 - Closing the mouth
 - Relaxed arms and hands
- Breastfeed or bottle-feed 8–12 times per day.
- Burp your baby during natural feeding breaks.
- Having 5–8 wet diapers and 3–4 stools each day shows your baby is eating well.

If Breastfeeding

- Continue to take your prenatal vitamins.
- When breastfeeding is going well (usually at 4–6 weeks), you can offer your baby a bottle or pacifier.

If Formula Feeding

- Always prepare, heat, and store formula safely. If you need help, ask us.
- Feed your baby 2 oz every 2–3 hours. If your baby is still hungry, you can feed more.
- Hold your baby so you can look at each other.
- Do not prop the bottle.

What to Expect at Your Baby's 2 Month Visit

We will talk about

- Taking care of yourself and your family
- Sleep and crib safety
- Keeping your home safe for your baby
- Getting back to work or school and finding child care
- Feeding your baby

Poison Help: 1-800-222-1222

Child safety seat inspection:
1-866-SEATCHECK; seatcheck.org



American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

The recommendations in this publication do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate. Original document included as part of Bright Futures Tool and Resource Kit. Copyright © 2010 American Academy of Pediatrics, Updated 10/11. All Rights Reserved. The American Academy of Pediatrics does not review or endorse any modifications made to this document and in no event shall the AAP be liable for any such changes.

A Parent's Guide to Safe Sleep

Helping you to reduce the risk of SIDS

DID YOU KNOW?

- About one in five sudden infant death syndrome (SIDS) deaths occur while an infant is in the care of someone other than a parent. Many of these deaths occur when babies who are used to sleeping on their backs at home are then placed to sleep on their tummies by another caregiver. We call this "unaccustomed tummy sleeping."
- Unaccustomed tummy sleeping increases the risk of SIDS. Babies who are used to sleeping on their backs and are placed to sleep on their tummies are 18 times more likely to die from SIDS.

You can reduce your baby's risk of dying from SIDS by talking to those who care for your baby, including child care providers, babysitters, family, and friends, about placing your baby to sleep on his back during naps and at night.

WHO IS AT RISK FOR SIDS?

- SIDS is the leading cause of death for infants between 1 month and 12 months of age.
- SIDS is most common among infants that are 1-4 months old. However, babies can die from SIDS until they are 1 year old.

KNOW THE TRUTH... SIDS IS NOT CAUSED BY:

- Immunizations
- Vomiting or choking

WHAT CAN I DO BEFORE MY BABY IS BORN TO REDUCE THE RISK OF SIDS?

Take care of yourself during pregnancy and after the birth of your baby. During pregnancy, before you even give birth, you can reduce the risk of your baby dying from SIDS! **Don't smoke or expose yourself to others' smoke while you are pregnant and after the baby is born. Alcohol and drug use can also increase your baby's risk for SIDS.** Be sure to visit a physician for regular prenatal checkups to reduce your risk of having a low birth weight or premature baby.

MORE WAYS TO PROTECT YOUR BABY

Do your best to follow the guidelines on these pages. This way, you will know that you are doing all that you can to keep your baby healthy and safe.

- Breastfeed your baby. Experts recommend that mothers feed their children human milk for as long and as much as possible, and for at least the first 6 months of life, if possible.
- It is important for your baby to be up to date on her immunizations and well-baby check-ups.

WHERE IS THE SAFEST PLACE FOR MY BABY TO SLEEP?

The safest place for your baby to sleep is in the room where you sleep, but not in your bed. Place the baby's crib or bassinet near your bed (within arm's reach). This makes it easier to breastfeed and to bond with your baby.

The crib or bassinet should be free from toys, soft bedding, blankets, and pillows. (See picture on next page.)

TALK ABOUT SAFE SLEEP PRACTICES WITH EVERYONE WHO CARES FOR YOUR BABY!

When looking for someone to take care of your baby, including a child care provider, a family member, or a friend, make sure that you talk with this person about safe sleep practices. Bring this fact sheet along to help, if needed. If a caregiver does not know the best safe sleep practices, respectfully try to teach the caregiver what you have learned about safe sleep practices and the importance of following these rules when caring for infants. Before leaving your baby with anyone, be sure that person agrees that the safe sleep practices explained in this brochure will be followed all of the time.



Face up to wake up – healthy babies sleep safest on their backs.



Do not place pillows, quilts, toys, or anything in the crib.



Supervised, daily tummy time during play is important to baby's healthy development.

WHAT ELSE CAN I DO TO REDUCE MY BABY'S RISK?

Follow these easy and free steps to help you reduce your baby's risk of dying from SIDS.

SAFE SLEEP PRACTICES

- Always place babies to sleep on their backs during naps and at nighttime. Because babies sleeping on their sides are more likely to accidentally roll onto their stomach, the side position is just as dangerous as the stomach position.
- Avoid letting the baby get too hot. The baby could be too hot if you notice sweating, damp hair, flushed cheeks, heat rash, and rapid breathing. Dress the baby lightly for sleep. Set the room temperature in a range that is comfortable for a lightly clothed adult.
- Consider using a pacifier at nap time and bed time. The pacifier should not have cords or clips that might be a strangulation risk.

SAFE SLEEP ENVIRONMENT

- Place your baby on a firm mattress, covered by a fitted sheet that meets current safety standards. For more about crib safety standards, visit the Consumer Product Safety Commission's Web site at <http://www.cpsc.gov>.
- Place the crib in an area that is always smoke free.
- Don't place babies to sleep on adult beds, chairs, sofas, waterbeds, pillows, or cushions.
- Toys and other soft bedding, including fluffy blankets, comforters, pillows, stuffed animals, bumper pads, and wedges should not be placed in the crib with the baby. Loose bedding, such as sheets and blankets, should not be used as these items can impair the infant's ability to breathe if they are close to his face. Sleep clothing, such as sleepers, sleep sacks, and wearable blankets are better alternatives to blankets.

IS IT EVER SAFE TO HAVE BABIES ON THEIR TUMMIES?

Yes! You should talk to your child care provider about making tummy time a part of your baby's daily activities. Your baby needs plenty of tummy time while supervised and awake to help build strong neck and shoulder muscles. Remember to make sure that your baby is having tummy time at home with you.

TUMMY TO PLAY AND BACK TO SLEEP

- Place babies to sleep on their backs to reduce the risk of SIDS. Side sleeping is not as safe as back sleeping and is not advised. Babies sleep comfortably on their backs, and no special equipment or extra money is needed.
- "Tummy time" is playtime when infants are awake and placed on their tummies while someone is watching them. Have tummy time to allow babies to develop normally.

WHAT CAN I DO TO HELP SPREAD THE WORD ABOUT BACK TO SLEEP?

- Be aware of safe sleep practices and how they can be made a part of our everyday lives.
- When shopping in stores with crib displays that show heavy quilts, pillows, and stuffed animals, talk to the manager about safe sleep, and ask them not to display cribs in this way.
- Monitor the media. When you see an ad or a picture in the paper that shows a baby sleeping on her tummy, write a letter to the editor.
- If you know teenagers who take care of babies, talk with them. They may need help with following the proper safe sleep practices.
- Set a good example – realize that you may not have slept on your back as a baby, but we now know that this is the safest way for babies to sleep. When placing babies to sleep, be sure to always place them on their backs.

If you have questions about safe sleep practices please contact Healthy Child Care America at the American Academy of Pediatrics at childcare@aap.org or 888/227-5409. Remember, if you have a question about the health and safety of your child, talk to your baby's doctor.

RESOURCES:

- American Academy of Pediatrics
<http://www.aappolicy.org>
SIDS and Other Sleep-Related Infant Deaths: Expansion of Recommendations for a Safe Infant Sleeping Environment
<http://aappolicy.aappublications.org/cgi/rep rint/pediatrics;128/5/e1341.pdf>
- Healthy Child Care America
<http://www.healthychildcare.org>
- National Resource Center for Health and Safety in Child Care and Early Education
<http://nrc.uchsc.edu>
- Healthy Kids, Healthy Care: A Parent Friendly Tool on Health and Safety Issues in Child Care <http://www.healthykids.us>
- National Institute for Child and Human Development Back to Sleep Campaign (Order free educational materials)
<http://www.nichd.nih.gov/sids/sids.cfm>
- First Candle/SIDS Alliance
<http://www.firstcandle.org>
- Association of SIDS and Infant Mortality Programs <http://www.asip1.org>
- CJ Foundation for SIDS
<http://www.cjsids.com>
- National SIDS and Infant Death Resource Center <http://www.sidscenter.org/>
- The Juvenile Products Manufacturers Association <http://www.jpma.org/>

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Revised 2012